



British  
Geological  
Survey



MONTHLY MAGNETIC BULLETIN

# Hartland Observatory

June 2023



# Hartland Observatory magnetic data

## 1. Introduction

Hartland Observatory is one of three geomagnetic observatories in the UK operated and maintained by the British Geological Survey (BGS).

This bulletin is published to provide rapid access to the provisional geomagnetic observatory results. The information is freely available for personal, academic, educational and non-commercial research or use. Magnetic observatory data are presented as a series of plots of one-minute, hourly and daily values, followed by tabulations of monthly values, reports of rapid variations and geomagnetic activity indices. The operation of the observatory and presentation of data are described in the rest of this section.

Enquiries about the data should be addressed to:

Geomagnetism Team  
British Geological Survey  
The Lyell Centre  
Research Avenue South  
Edinburgh EH14 4AP  
Scotland, UK

Tel: +44 (0) 131 667 1000  
Email: [enquiries@bgs.ac.uk](mailto:enquiries@bgs.ac.uk)  
Internet: [geomag.bgs.ac.uk](http://geomag.bgs.ac.uk)

## 2. Position

The observatory is situated on the NW boundary of the village of Hartland in North Devon. The observatory co-ordinates are:

*Geographic:* 50° 59'42.0"N 355° 30'57.6"E  
*Geomagnetic:* 53° 34'40.8"N 079° 45'54.0"E  
*Height above mean sea level:* 95 m

The geographical coordinates are measured by a handheld GPS device, which uses WGS84 as the reference coordinate system. The height above MSL is determined from the best available contour maps. The geomagnetic co-ordinates are approximations, calculated using the 13th generation International Geomagnetic Reference Field (IGRF) at epoch 2021.5. On-line access to models (including IGRF), charts and navigational data are available at [geomag.bgs.ac.uk/data\\_service/models\\_compass/home](http://geomag.bgs.ac.uk/data_service/models_compass/home)

## 3. The observatory operation

### 3.1 GDAS

The observatory operates under the control of the Geomagnetic Data Acquisition System (GDAS), which was developed by BGS staff, installed in 2002, and became fully operational in January 2003. The data acquisition software, running on QNX operated computers, controls the data logging and the communications.

There are two sets of sensors used for making magnetic measurements. A tri-axial linear-core fluxgate magnetometer, manufactured by the Danish Meteorological Institute, is used to measure the variations in the horizontal ( $H$ ) and vertical ( $Z$ ) components of the field. The third sensor is oriented perpendicular to these, and measures variations, which are proportional to the changes in declination ( $D$ ). Measurements are made at a rate of 1 Hz.

In addition to the fluxgate sensors there is a proton precession magnetometer (PPM) making measurements of the absolute total field intensity ( $F$ ) at a rate of 0.1Hz.

The raw unfiltered data are retrieved automatically via Internet connections to the BGS office in Edinburgh in near real-time. The fluxgate data are filtered to produce one-minute values using a 61-point cosine filter and the total field intensity samples are filtered using a 7-point cosine filter. The one-minute values provide input for various data products, available on-line at [geomag.bgs.ac.uk/data\\_service/home](http://geomag.bgs.ac.uk/data_service/home)

### 3.2 Back-up systems

There are two other fully independent identical systems, GDAS 2 and GDAS 3, operating at the observatory. The data from these are also processed in near real-time and used for quality control purposes. They are also used to fill any gaps or replace any corrupt values in the primary system, GDAS 1.

### 3.3 Absolute observations

The GDAS fluxgate magnetometers accurately measure variations in the components of the geomagnetic field, but not the absolute magnitudes. Two sets of absolute measurements of the field are made manually once per week. A fluxgate sensor mounted on a theodolite is used to determine  $D$  and inclination ( $I$ ); the GDAS PPM measurements, with a site difference correction applied, are used for  $F$ . The absolute observations are used in conjunction with the GDAS variometer measurements to produce a continuous record of the absolute values of the geomagnetic field elements as if they had been measured at the observatory reference pillar.

## 4. Observatory results

The data presented in the bulletin are in the form of plots and tabulations described in the following sections.

### 4.1 Absolute observations

The absolute observation measurements made during the month are tabulated. Also included are the corresponding baseline values, which are the differences between the absolute measurements and the variometer measurements of  $D$ ,  $H$  and  $Z$  (in the sense absolute-variometer). These are also plotted (markers) along with the derived preliminary daily baseline values (line) throughout the year. Daily mean differences between the measured absolute  $F$  and the  $F$  computed from the baseline corrected  $H$  and  $Z$  values are plotted in the fourth panel (in the sense measured-derived). The bottom panel shows the daily mean temperature in the fluxgate chamber.

## 4.2 Summary magnetograms

Small-scale magnetograms are plotted which allow the month's data to be viewed at a glance. They are plotted 16 days to a page and show the one-minute variations in  $D$ ,  $H$  and  $Z$ . The scales are shown on the right-hand side of the page. On disturbed days the scales are multiplied by a factor, which is indicated above the panel for that day. The variations are centred on the monthly mean value, shown on the left side of the page.

## 4.3 Magnetograms

The daily magnetograms are plotted using one-minute values of  $D$ ,  $H$  and  $Z$  from the fluxgate sensors, with any gaps filled using back-up data. The magnetograms are plotted to a variable scale; scale bars are shown to the right of each plot. The absolute level (the monthly mean value) is indicated on the left side of the plots.

## 4.4 Hourly mean value plots

Hourly mean values of  $D$ ,  $H$  and  $Z$  for the past 12 months are plotted in 27-day segments corresponding to the Bartels solar rotation number. Magnetic disturbances associated with active regions and/or coronal holes on the Sun may recur after 27 days: the same is true for geomagnetically quiet intervals. Plotting the data in this way highlights this recurrence. Diurnal variations are also clear in these plots and the amplitude changes throughout the year highlight the seasonal changes. Longer term secular variation is also illustrated.

Full lists of the UK observatory hourly mean values from 1983 to the present day are available at [geomag.bgs.ac.uk/data\\_service/data/obs\\_data/hourly\\_means](http://geomag.bgs.ac.uk/data_service/data/obs_data/hourly_means)

## 4.5 Daily and monthly mean values

Daily mean values of  $D$ ,  $H$ ,  $Z$  and  $F$  are plotted throughout the year. In addition, a table of monthly mean values of all the geomagnetic elements is provided. These values depend on accurate specification of the fluxgate sensor baselines. It is anticipated that these provisional values will not be altered by more than a few nT or tenths of arcminutes before being made definitive at the end of the year.

## 4.6 Rapid variations

Charged particles stream from the Sun in the solar wind. The solar wind interacts with the geomagnetic field to create a cavity, the magnetosphere, in which the field is confined. When a region of enhanced velocity and/or density in the solar wind arrives at the dayside boundary of the magnetosphere (at about 10 earth radii) the boundary is pushed towards the Earth. Currents set up on the boundary of the magnetosphere can cause an abrupt change in the geomagnetic field measured on the ground and this is recorded on observatory magnetograms as a sudden impulse ( $si$ ). If, following an  $si$ , there is a change in the rhythm of activity, the  $si$  is termed a storm sudden commencement ( $ssc$ ). A classical magnetic storm exhibiting initial, main and recovery phases (shown by, for instance, the  $Dst$  ring current index) can often occur after a  $ssc$ , in which case the start of the storm is taken as the time of the  $ssc$ .

Solar flares, seen at optical wavelengths as a sudden brightening of a small region of the Sun's surface, are also responsible for increased X-ray emissions. These X-rays cause increased ionisation in the ionosphere, which leads to absorption of short-wave radio signals. A solar flare effect ( $sfe$ ), or "crochet", may be observed on a magnetogram during geomagnetically quiet times. It is a relatively short-term change (tens of minutes) to the normal diurnal variation and can vary in size (tens of nT) depending on local time (LT), geomagnetic latitude and solar zenith angle.

## 4.7 Local geomagnetic activity indices

*The Observatory K index.* This summarises geomagnetic activity at an observatory by assigning a code, an integer in the range 0 to 9, to each 3-hour Universal Time (UT) interval. The index for each 3-hour UT interval is determined from the maximum range in  $H$  or  $D$

(scaled in nT), with allowance made for the regular (undisturbed) diurnal variation. The conversion from range to an index value is made using a quasi-logarithmic scale, with the scale values dependent on the geomagnetic latitude of the observatory. The lower bounds (in nT) for the classification of each period at Hartland are:

0	1	2	3	4	5	6	7	8	9
0	5	10	20	40	70	120	200	330	500

The *K* index retains the LT and seasonal dependence of activity associated with the position of the observatory. The 3-hourly *K* indices for the month are tabulated and also plotted as a histogram. All UK observatory *K* indices are available at [geomag.bgs.ac.uk/data\\_service/data/magnetic\\_indices/k\\_indices](http://geomag.bgs.ac.uk/data_service/data/magnetic_indices/k_indices)

#### 4.8 Global geomagnetic activity indices

*The aa index.* A number of 3-hour geomagnetic indices are computed by combining *K* indices from networks of observatories to characterise global activity levels and to eliminate LT and seasonal effects. The simplest of these is the *aa* index, computed using the *K* indices from two approximately antipodal observatories: Hartland in the UK and Canberra in Australia. The *aa* index is calculated from linearisations of the Hartland and Canberra *K* indices, and has units of nT. The 3-hourly *aa* indices are tabulated along with the daily mean value of *aa* (denoted *Aa*), the mean values of *aa* for the intervals 00-12UT (*Aa<sub>am</sub>*) and 12-24UT (*Aa<sub>pm</sub>*) and the monthly mean value. The 3-hourly *aa* indices for the month are also plotted as a histogram.

Although the *aa* index is based on data from only two observatories, provided averages over 12 hours or longer are used, the index is strongly correlated with the *ap* and *am* indices, which are derived using data from more extensive observatory networks.

The *aa* indices listed in this bulletin are available at [geomag.bgs.ac.uk/data\\_service/data/magnetic\\_indices/aaindex](http://geomag.bgs.ac.uk/data_service/data/magnetic_indices/aaindex) as well as the full data set from 1868.

Definitive *aa* are published by the International Service for Geomagnetic Indices, EOST, 5 rue René Descartes, 67084 Strasbourg Cedex, France.

## 5. Conditions of use

The data presented in this bulletin are provided for personal, academic, educational, non-commercial research or other non-commercial use and are not for sale or distribution to third parties without written permission from BGS.

Reproduction of any part of this bulletin should be accompanied by the statement: 'Reproduced with the permission of the British Geological Survey ©UKRI. All rights Reserved'. Publications making use of the data should include an acknowledgment statement of the form: 'The results presented in this paper rely on the data collected at Hartland magnetic observatory, operated by the British Geological Survey.'


Commercial users can contact the geomagnetism team for information on the range of applications and services offered. Full contact details are available at [geomag.bgs.ac.uk/contactus/staff](http://geomag.bgs.ac.uk/contactus/staff)

# Hartland Observatory

## Absolute observations

Date	Day Number	Declination			Inclination		Total field		Horizontal intensity		Vertical intensity		Observer
		Time (UT)	Absolute (°)	Baseline (°)	Time (UT)	Absolute (°)	Site difference (nT)	Absolute corrected (nT)	Absolute (nT)	Baseline (nT)	Absolute (nT)	Baseline (nT)	
05-Jun-23	156	10:49	-0.8134	-0.8733	10:57	65.9862	17.4	48765.4	19845.4	99999.0	44544.6	99999.0	BC
05-Jun-23	156	11:02	-0.8226	-0.8683	11:09	65.9847	17.4	48765.4	19846.6	99999.0	44544.1	99999.0	BC
21-Jun-23	172	09:28	-0.7137	-0.8700	09:35	66.0126	17.4	48780.2	19830.9	19874.2	44567.3	44565.3	BC
21-Jun-23	172	09:46	-0.7340	-0.8733	09:54	66.0117	17.4	48777.5	19830.5	99999.0	44564.5	99999.0	BC
23-Jun-23	174	08:06	-0.6658	-0.8700	08:14	66.0190	17.4	48785.8	19828.2	19874.3	44574.6	44565.2	BC
23-Jun-23	174	08:19	-0.6698	-0.8717	08:27	66.0185	17.4	48784.1	19827.8	19874.5	44572.9	44565.1	BC
26-Jun-23	177	10:10	-0.7766	-0.8700	10:17	66.0228	17.4	48773.5	19820.3	99999.0	44564.7	99999.0	BC
26-Jun-23	177	10:22	-0.7785	-0.8717	10:30	66.0175	17.4	48775.3	19825.1	19873.9	44564.5	44565.4	BC
28-Jun-23	179	07:08	-0.7087	-0.8717	07:16	65.9935	17.4	48797.1	19852.6	19873.1	44576.2	44565.6	BC
28-Jun-23	179	07:22	-0.7040	-0.8700	07:30	65.9942	17.4	48795.3	19851.3	19874.2	44574.7	44565.2	BC


# Hartland 2023




# Hartland

# June

# 2023







Hartland


June

2023

Date: 01-06-2023


# Hartland

Day number: 152



Date: 02-06-2023


Day number: 153



Date: 03-06-2023


# Hartland

Day number: 154



Date: 04-06-2023


Day number: 155



Date: 05-06-2023


# Hartland

Day number: 156



Date: 06-06-2023


Day number: 157



Date: 07-06-2023


# Hartland

Day number: 158



Date: 08-06-2023

Day number: 159




Hour (UT)

Date: 09-06-2023


# Hartland

Day number: 160



Date: 10-06-2023


Day number: 161



Date: 11-06-2023


# Hartland

Day number: 162



Date: 12-06-2023

Day number: 163




Hour (UT)

Date: 13-06-2023


# Hartland

Day number: 164



Date: 14-06-2023

Day number: 165






Date: 15-06-2023


# Hartland

Day number: 166



Date: 16-06-2023


Day number: 167



Date: 17-06-2023


# Hartland

Day number: 168



Date: 18-06-2023

Day number: 169




Hour (UT)

Date: 19-06-2023


# Hartland

Day number: 170



Date: 20-06-2023


Day number: 171



Date: 21-06-2023


# Hartland

Day number: 172



Date: 22-06-2023


Day number: 173



Date: 23-06-2023


# Hartland

Day number: 174



Date: 24-06-2023


Day number: 175



Date: 25-06-2023


# Hartland

Day number: 176



Date: 26-06-2023

Day number: 177




Hour (UT)

Date: 27-06-2023


# Hartland

Day number: 178



Date: 28-06-2023


Day number: 179



Date: 29-06-2023


# Hartland

Day number: 180




Date: 30-06-2023

Day number: 181






# Hartland Observatory: Declination (degrees)




Hourly Mean Values Plotted By Bartels Solar Rotation Number

# Hartland Observatory: Horizontal Intensity (nT)




# Hartland Observatory: Vertical Intensity (nT)



Hourly Mean Values Plotted By Bartels Solar Rotation Number

# Hartland Observatory 2023



## Monthly mean values for Hartland Observatory 2023

Month	<i>D</i>	<i>H</i>	<i>I</i>	<i>X</i>	<i>Y</i>	<i>Z</i>	<i>F</i>
January	-0° 52.2′	19838 nT	66° 0.2′	19835 nT	-301 nT	44564 nT	48780 nT
February	-0° 51.0′	19833 nT	66° 0.6′	19831 nT	-294 nT	44566 nT	48780 nT
March	-0° 49.7′	19832 nT	66° 0.7′	19830 nT	-287 nT	44570 nT	48783 nT
April	-0° 49.1′	19840 nT	66° 0.2′	19838 nT	-283 nT	44568 nT	48784 nT
May	-0° 48.1′	19847 nT	65° 59.8′	19845 nT	-278 nT	44571 nT	48790 nT
June	-0° 47.6′	19853 nT	65° 59.4′	19851 nT	-275 nT	44570 nT	48792 nT

### Note:

i. The values shown here are provisional.

## Hartland rapid variations

### SIs and SSCs

Date	Time (UT)	Type	Quality	H (nT)	D (min)	Z (nT)
None						

#### Notes:

An asterisk (\*) indicates that the principal impulse was preceded by a smaller reversed impulse.

The quality of the event is classified as follows:

- A = very distinct
- B = fair, ordinary, but unmistakable
- C = doubtful

The amplitudes given are for the first chief movement of the event.

### SFEs


Date	Start	Universal Time Maximum	End	H (nT)	D (min)	Z (nT)
None						

#### Note:

The amplitudes given are for the first chief movement of the event.

# Indices of geomagnetic activity

Day	K - INDICES FOR THREE-HOUR INTERVAL							
	00-03	03-06	06-09	09-12	12-15	15-18	18-21	21-24
1	4	3	4	1	3	4	3	2
2	3	1	1	0	1	1	1	2
3	1	1	1	0	1	0	1	1
4	2	1	2	2	2	2	4	3
5	0	0	1	1	0	1	2	2
6	3	2	1	1	2	0	1	0
7	0	1	1	1	1	1	1	2
8	1	1	1	1	2	2	2	0
9	1	1	0	0	0	2	2	1
10	1	0	1	1	2	2	2	1
11	2	3	2	1	2	3	2	1
12	1	2	2	1	3	2	2	1
13	1	1	1	1	2	3	1	1
14	3	1	1	1	0	1	0	0
15	1	1	2	1	4	5	4	5
16	4	5	3	4	4	4	2	3
17	2	3	2	2	2	4	2	2
18	3	2	1	1	2	2	4	2
19	2	1	2	1	3	4	3	4
20	3	2	1	2	2	3	2	2
21	2	2	1	0	2	1	2	3
22	1	2	3	3	3	3	4	3
23	3	2	1	1	2	3	3	2
24	1	3	4	2	3	4	4	4
25	4	2	2	1	2	4	4	3
26	2	2	2	2	3	4	2	3
27	2	2	2	2	2	3	2	2
28	2	3	1	2	2	2	2	2
29	3	3	4	3	1	4	3	3
30	3	2	1	1	4	2	0	0



## The aa Index

Date	Day	3-hourly aa-indices								$Aa_{am}$	$Aa_{pm}$	$Aa$
01/06/2023	152	45	32	59	20	24	45	20	12	39.1	25.3	32.2
02/06/2023	153	17	8	12	2	5	5	8	9	9.9	6.8	8.3
03/06/2023	154	8	8	5	5	5	2	5	8	6.5	5.1	5.8
04/06/2023	155	12	5	16	24	16	12	37	20	14.3	21.3	17.8
05/06/2023	156	5	5	8	12	2	5	12	16	7.5	8.9	8.2
06/06/2023	157	24	24	8	12	12	2	5	2	17	5.5	11.2
07/06/2023	158	5	5	5	5	5	5	5	16	5.1	7.9	6.5
08/06/2023	159	12	12	8	12	16	9	9	2	11	9.2	10.1
09/06/2023	160	5	5	2	2	2	12	16	12	3.8	10.6	7.2
10/06/2023	161	8	2	5	8	12	9	12	5	5.8	9.5	7.7
11/06/2023	162	16	32	16	8	12	20	12	8	18	12.9	15.4
12/06/2023	163	8	9	12	5	20	12	12	5	8.5	12.2	10.3
13/06/2023	164	5	8	8	8	12	20	5	5	7.1	10.5	8.8
14/06/2023	165	20	8	5	5	2	5	2	2	9.5	3.1	6.3
15/06/2023	166	8	12	16	12	45	67	81	80	12	68.2	40.1
16/06/2023	167	45	80	46	81	59	45	24	20	63	37.1	50
17/06/2023	168	16	20	16	16	16	37	16	12	17	20.3	18.7
18/06/2023	169	24	12	8	20	9	24	45	12	16	22.7	19.3
19/06/2023	170	12	8	12	12	32	45	32	37	10.9	36.7	23.8
20/06/2023	171	20	12	8	12	24	24	12	12	12.9	18	15.4
21/06/2023	172	12	16	8	5	12	5	9	24	10.2	12.6	11.4
22/06/2023	173	8	12	32	24	32	24	33	24	19	28.3	23.6
23/06/2023	174	20	12	12	20	16	32	32	16	16	24.1	20
24/06/2023	175	8	32	45	16	32	37	37	45	25.4	38	31.7
25/06/2023	176	45	24	16	8	24	45	37	24	23.4	32.7	28
26/06/2023	177	12	12	16	16	24	45	12	24	14	26.3	20.2
27/06/2023	178	16	12	16	24	16	20	12	12	17.1	14.9	16
28/06/2023	179	12	20	8	16	12	9	12	12	13.9	11.2	12.6
29/06/2023	180	20	32	45	46	12	45	32	24	35.8	28.4	32.1
30/06/2023	181	24	12	8	20	45	12	2	2	16	15.6	15.8
Monthly Mean Value											17.8	

### Notes

- i. The units of the aa index are nT.
- ii. The 3-hour aa values are rounded to the nearest integer. Where  $aa = *.5$ , aa is rounded down.
- iii. Daily values ( $Aa_{am}$ ,  $Aa_{pm}$  and  $Aa$ ) are computed from aa values of original resolution.
- iv. The monthly mean value is computed from the daily mean values,  $Aa$ .
- v. Definitive aa indices are derived and published by the International Service for Geomagnetic Indices.

