


**BRITISH GEOLOGICAL SURVEY**

**Jim Carrigan  
Observatory  
Prudhoe Bay  
Monthly  
Magnetic  
Bulletin  
August 2009  
09/08/JC**



**British Geological Survey**

NATURAL ENVIRONMENT RESEARCH COUNCIL

# 1. JIM CARRIGAN OBSERVATORY MAGNETIC DATA

## 1. Introduction

Jim Carrigan Observatory is the fourth overseas geomagnetic observatory established by the British Geological Survey (BGS). The installation is a joint project between BGS and Sperry Drilling Services (SDS) in support of directional drilling programmes. SDS has operated a prototype station since 1997 which was upgraded by the BGS to a standard high-quality observatory in October 2003.

This bulletin is published to meet the needs of users of geomagnetic data. Magnetic observatory data is presented as a series of plots of one-minute, hourly and daily values, followed by a tabulation of monthly values. The operation of the observatory and presentation of data are described in the rest of this section.

Enquiries about the data should be addressed to:

National Geomagnetic Service  
British Geological Survey  
Murchison House, West Mains Road  
Edinburgh EH9 3LA  
Scotland, UK

Tel: +44 (0) 131 667 1000  
Fax: +44 (0) 131 668 0265  
E-mail: orba@bgs.ac.uk  
Internet: www.geomag.bgs.ac.uk

## 2. Position

Jim Carrigan Observatory is situated at T-Pad, a man-made gravel bed close to the drilling sites at Prudhoe Bay, Alaska, USA.

The observatory co-ordinates are:-

*Geographic:* 70° 21.36'N 211° 12.06'E  
*Geomagnetic:* 70° 04.2'N 254° 00.36'E  
*Height above mean sea level:* 10m (approx)

The geomagnetic co-ordinates are calculated using the 10<sup>th</sup> generation International Geomagnetic Reference Field (IGRF) at epoch 2009.5.

## 3. The Observatory Operation

### 3.1 GDAS

The observatory operates under the control of the Geomagnetic Data Acquisition System (GDAS), which was developed by BGS and became fully operational from 27<sup>th</sup> October 2003. The system operates under the control of data acquisition software running on QNX computers, which controls the data logging and communications.

There are two sets of sensors used for making magnetic measurements. A triaxial linear-core fluxgate magnetometer, manufactured by the Danish Meteorological Institute, is used to measure the variations in the horizontal (*H*) and vertical (*Z*) components of the field. The third sensor is oriented perpendicular to these, and measures variations, which are proportional to the changes in declination (*D*). Measurements are made at a rate of 1 Hz.

In addition to the fluxgate sensors there is a proton precession magnetometer making measurements of the absolute total field intensity (*F*) at a rate of 0.1Hz.

The raw unfiltered data are retrieved automatically via internet connections to the BGS office in Edinburgh in near real-time. The fluxgate data are filtered to produce one-minute values using a 61-point cosine filter whilst the total field intensity samples are filtered using a 7-point cosine filter.

### 3.2 Absolute Observations

The GDAS fluxgate magnetometers accurately measure variations in the components of the geomagnetic field, but not the absolute magnitudes. Regular manual absolute measurements of the field are made throughout the year. A fluxgate sensor mounted on a theodolite is used to determine *D* and inclination (*I*); the GDAS PPM measurements, with a site difference correction applied, are used for *F*. The absolute observations are used in conjunction with the GDAS variometer measurements to produce a continuous record of the absolute values of the geomagnetic field elements as if they had been measured at the observatory reference pillar.

## 4. Data Presentation

The data presented in the bulletin are in the form of plots and tabulations described in the following sections.

### 4.1 Absolute Observations

The absolute observation measurements made during the month are tabulated. Also included are the corresponding baseline values, which are the differences between the absolute measurements and the variometer measurements of *D*, *H* and *Z* (in the sense absolute–variometer). These are also plotted (markers) along with the derived preliminary daily baseline values (line) throughout the year. Daily mean differences between the measured absolute *F* and the *F* computed from the baseline corrected *H* and *Z* values are plotted in the fourth panel (in the sense measured–derived). The bottom panel shows the daily mean temperature in the fluxgate chamber.

#### **4.2 Summary magnetograms**

Small-scale magnetograms are plotted which allow the month's data to be viewed at a glance. They are plotted 16 days a page and show the variations in  $D$ ,  $H$  and  $Z$ . The scales are shown on the right-hand side of the page. On disturbed days the scales are multiplied by a factor, which is indicated above the panel for that day. The variations are centred on the monthly mean value, shown on the left side of the page.

#### **4.3 Magnetograms**

The daily magnetograms are plotted using one-minute values of  $D$ ,  $H$  and  $Z$  from the fluxgate sensors, with any gaps filled using back-up data. The magnetograms are plotted to a variable scale; scale bars are shown to the right of each plot. The absolute level (the monthly mean value) is indicated on the left side of the plots.

#### **4.4 Hourly Mean Value Plots**

Hourly mean values of  $D$ ,  $H$  and  $Z$  for the past 12 months are plotted in 27-day segments corresponding to the Bartels solar rotation number. Magnetic disturbances associated with active regions on the surface of the Sun may recur after 27 days: the same is true for geomagnetically quiet intervals. Plotting the data in this way highlights this recurrence, and also illustrates seasonal and diurnal variations throughout the year.


#### **4.5 Daily and Monthly Mean Values**

Daily mean values of  $D$ ,  $H$ ,  $Z$  and  $F$  are plotted throughout the year. In addition, a table of monthly mean values of all the geomagnetic elements is provided. These values depend on accurate specification of the fluxgate sensor baselines.






# Jim Carrigan Obs 2009




Jim Carrigan

August

2009






Jim Carrigan August 2009



Date: 01-08-2009


Jim Carrigan

Day number: 213



Date: 02-08-2009


Day number: 214



Date: 03-08-2009


Jim Carrigan

Day number: 215



Date: 04-08-2009


Day number: 216



Date: 05-08-2009


Jim Carrigan

Day number: 217



Date: 06-08-2009


Day number: 218



Date: 07-08-2009


Jim Carrigan

Day number: 219



Date: 08-08-2009


Day number: 220



Date: 09-08-2009


Jim Carrigan

Day number: 221



Date: 10-08-2009


Day number: 222



Date: 11-08-2009


Jim Carrigan

Day number: 223



Date: 12-08-2009


Day number: 224



Date: 13-08-2009


Jim Carrigan

Day number: 225



Date: 14-08-2009


Day number: 226



Date: 15-08-2009


Jim Carrigan

Day number: 227



Date: 16-08-2009

Day number: 228






Date: 17-08-2009


Jim Carrigan

Day number: 229



Date: 18-08-2009

Day number: 230




Hour (UT)

Date: 19-08-2009


Jim Carrigan

Day number: 231



Date: 20-08-2009


Day number: 232



Date: 21-08-2009


Jim Carrigan

Day number: 233



Date: 22-08-2009


Day number: 234



Date: 23-08-2009


Jim Carrigan

Day number: 235



Date: 24-08-2009


Day number: 236



Date: 25-08-2009


Jim Carrigan

Day number: 237



Date: 26-08-2009


Day number: 238



Date: 27-08-2009


Jim Carrigan

Day number: 239



Date: 28-08-2009


Day number: 240



Date: 29-08-2009


Jim Carrigan

Day number: 241



Date: 30-08-2009


Day number: 242



Date: 31-08-2009


Jim Carrigan

Day number: 243







# Jim Carrigan Observatory: Declination (degrees)




# Jim Carrigan Observatory: Horizontal Intensity (nT)



# Jim Carrigan Observatory: Vertical Intensity (nT)



# Jim Carrigan Observatory 2009



### Monthly Mean Values for Jim Carrigan Observatory 2009

Month	<i>D</i>	<i>H</i>	<i>I</i>	<i>X</i>	<i>Y</i>	<i>Z</i>	<i>F</i>
January	22° 45.9′	9018 nT	80° 59.6′	8316 nT	3490 nT	56893 nT	57603 nT
February	22° 43.5′	9019 nT	80° 59.6′	8319 nT	3484 nT	56896 nT	57606 nT
March	22° 40.6′	9019 nT	80° 59.6′	8322 nT	3477 nT	56898 nT	57608 nT
April	22° 38.1′	9020 nT	80° 59.5′	8325 nT	3471 nT	56898 nT	57608 nT
May	22° 34.7′	9028 nT	80° 59.0′	8336 nT	3467 nT	56898 nT	57610 nT
June	22° 31.9′	9035 nT	80° 58.6′	8345 nT	3462 nT	56897 nT	57610 nT
July	22° 28.8′	9025 nT	80° 59.3′	8339 nT	3451 nT	56904 nT	57615 nT
August	22° 26.3′	9021 nT	80° 59.5′	8338 nT	3443 nT	56905 nT	57616 nT

#### Note

i. The values shown here are provisional.