



# Monthly Geomagnetic Bulletin

September 2003 03/09/LE



# 1. LERWICK OBSERVATORY MAGNETIC DATA

## 1.1 Introduction

This bulletin is published to meet the needs of both commercial and academic users of geomagnetic data. Magnetic observatory data is presented as a series of plots of one-minute, hourly and daily values, followed by tabulations of monthly values, geomagnetic activity indices and reports of rapid variations. The operation of the observatory and presentation of data are described in the rest of this section.

Enquiries about the data should be addressed to:

National Geomagnetic Service  
British Geological Survey  
Murchison House, West Mains Road  
Edinburgh EH9 3LA  
Scotland, UK

Tel: +44 (0) 131 667 1000  
Fax: +44 (0) 131 668 4368  
E-mail: [s.reay@bgs.ac.uk](mailto:s.reay@bgs.ac.uk)  
Internet: [www.geomag.bgs.ac.uk](http://www.geomag.bgs.ac.uk)

## 1.2 Position

Lerwick Observatory, one of the three geomagnetic observatories operated and maintained in the UK by BGS, is situated on a ridge of high ground about 2.5 km to the SW of the port of Lerwick in Shetland. The observatory co-ordinates are:

*Geographic:* 60°08.0'N 358°49.0'E  
*Geomagnetic:* 62°00.0'N 88°59.0'E  
*Height above mean sea level:* 85 m

The geomagnetic co-ordinates are calculated using the 8th generation International Geomagnetic Reference Field at epoch 2003.5.

## 1.3 The Observatory Operation

### 1.3.1 GDAS

The observatory operates under the control of the Geomagnetic Data Acquisition System (GDAS), which was developed by BGS staff, installed in 2002, and became fully operational in January 2003. The system operates under the control of data acquisition software running on QNX computers, which control the data-logging and communications.

There are two sets of sensors used for making magnetic measurements. A triaxial linear-core fluxgate magnetometer, manufactured by the Danish Meteorological Institute, is used to measure the variations in the horizontal ( $H$ ) and vertical ( $Z$ ) components of the field. The third sensor is oriented perpendicular to these, and measures variations which are proportional to the changes in declination ( $D$ ). Measurements are made at a rate of 1 Hz.

In addition to the fluxgate sensors there is a proton precession magnetometer making measurements of the absolute total field intensity ( $F$ ) at a rate of 0.1Hz.

The raw unfiltered data are retrieved automatically via Internet connections to the BGS office in Edinburgh in near real-time. The fluxgate data are filtered to produce one-minute values using a 61-point cosine filter whilst the total field intensity samples are filtered using a 7-point cosine filter. These one-minute values are used to update the Geomagnetism Information and Forecast Service (GIFS), an on-line information system accessed via the World Wide Web at the address given in Section 1.1. GIFS also provides information on geomagnetic and solar activity.

### 1.3.2 Back-up Systems

There are two other fully independent identical systems, GDAS 2 and GDAS 3, operating at the observatory. The data from these are also processed in near real-time and used for quality control purposes. They can be also be used to fill any gaps or replace any corrupt values in the primary system, GDAS 1.

## 1.4 Data Presentation

The data presented in the bulletin are in the form of plots and tabulations described in the following sections.

### 1.4.1 Summary magnetograms

Small-scale magnetograms are plotted which allow the month's data to be viewed at a glance. They are plotted 16 days a page and show the variations in  $D$ ,  $H$  and  $Z$ . The scales are shown on the right-hand side of the page. On disturbed days the scales are multiplied by a factor, which is indicated above the panel for that day. The variations are centred on the monthly mean value, shown on the left side of the page.

### 1.4.2 Magnetograms

The daily magnetograms are plotted using one-minute values of  $D$ ,  $H$  and  $Z$  from the fluxgate sensors, with any gaps filled using back-up data. The magnetograms are plotted to a variable scale; scale bars are shown to the right of each plot. The absolute level (the monthly mean value) is indicated on the left side of the plots.

### 1.4.3 Hourly Mean Value Plots

Hourly mean values of  $D$ ,  $H$  and  $Z$  for the past 12 months are plotted in 27-day segments corresponding to the Bartels solar rotation number. Magnetic disturbances associated with active regions on the surface of the Sun may recur after 27 days: the same is true for geomagnetically quiet intervals. Plotting the data in this way highlights this recurrence, and also illustrates seasonal and diurnal variations throughout the year.

### 1.4.4 Daily and Monthly Mean Values

Daily mean values of  $D$ ,  $H$ ,  $Z$  and  $F$  are plotted throughout the year. In addition, a table of monthly mean values of all the geomagnetic elements is provided. These values depend on accurate specification of the fluxgate sensor baselines. Provisional and definitive values are indicated in the table as **P** or **D** respectively. It is anticipated that provisional values will not be altered by more than a few nT or tenths of arcminutes before being made definitive.

### 1.4.5 Geomagnetic activity indices

*The Observatory K index.* This summarises geomagnetic activity at an observatory by assigning a code, an integer in the range 0 to 9, to each 3-hour Universal Time (UT) interval. The index for each 3-hour UT interval is determined from the ranges in  $H$  and in  $D$  (scaled in nT), with allowance made for the regular (undisturbed) diurnal variation. The conversion from range to an index value is made using a quasi-logarithmic scale, with the scale values dependent on the geomagnetic latitude of the observatory. The  $K$  index retains the local time (LT) and seasonal dependence of activity associated with the position of the observatory.

*The provisional aa index.* A number of 3-hour geomagnetic indices are computed by combining  $K$  indices from networks of observatories to characterise global activity levels and to eliminate LT and seasonal effects. The simplest of these is the  $aa$  index, computed using the  $K$  indices from two approximately antipodal observatories: Hartland in the UK and Canberra in Australia. The  $aa$  index is calculated from linearisations of the Hartland and Canberra  $K$  indices, and has units of nT. The daily

mean value of  $aa$  (denoted  $Aa$ ), the mean values of  $aa$  for the intervals 00-12UT and 12-24UT and the daily mean values for Hartland alone ( $Aa_n$ ) and Canberra alone ( $Aa_s$ ) are tabulated.

Although the  $aa$  index is based on data from only two observatories, provided averages over 12 hours or longer are used, the index is strongly correlated with the  $ap$  and  $am$  indices, which are derived using data from more extensive observatory networks.

The  $aa$  indices listed in this publication are provisional only; the definitive values are published by the International Service for Geomagnetic Indices, CRPE/CNET - CNRS, 4 Avenue de Neptune, F-94107 Saint Maur Cedex, France.

### 1.4.6 Rapid Variations


Charged particles stream from the Sun in the solar wind. The solar wind interacts with the geomagnetic field to create a cavity, the magnetosphere, in which the field is confined. When a region of enhanced velocity and/or density in the solar wind arrives at the dayside boundary of the magnetosphere (at about 10 earth radii) the boundary is pushed towards the Earth. Currents set up on the boundary of the magnetosphere can cause an abrupt change in the geomagnetic field measured on the ground and this is recorded on observatory magnetograms as a Sudden Impulse (SI). If, following an SI, there is a change in the rhythm of activity, the SI is termed a Storm Sudden Commencement (SSC). A classical magnetic storm exhibiting initial, main and recovery phases (shown by, for instance, the  $Dst$  ring current index) can often occur after a SSC, in which case the start of the storm is taken as the time of the SSC.


Solar flares, seen at optical wavelengths as a sudden brightening of a small region of the Sun's surface, are also responsible for increased X-ray emissions. The X-rays cause increased ionisation in the ionosphere, which leads to absorption of short-wave radio signals. On an observatory magnetogram a Solar Flare Effect (SFE), or "crochet" may be observed. This is an enhancement to the diurnal variation of the order of 10 nT, lasting about an hour.

Lerwick

September

2003





Lerwick


September

2003

Date: 01-09-2003


# Lerwick

Day number: 244



Date: 02-09-2003


Day number: 245



Date: 03-09-2003


# Lerwick

Day number: 246



Date: 04-09-2003


Day number: 247



Date: 05-09-2003


# Lerwick

Day number: 248



Date: 06-09-2003

Day number: 249






Date: 07-09-2003


# Lerwick

Day number: 250



Date: 08-09-2003


Day number: 251



Date: 09-09-2003


# Lerwick

Day number: 252




Date: 10-09-2003


Day number: 253



**Date: 11-09-2003** **Lerwick** **Day number: 254**




**Date: 12-09-2003** **Day number: 255**



Date: 13-09-2003


# Lerwick

Day number: 256



Date: 14-09-2003


Day number: 257



Date: 15-09-2003


# Lerwick

Day number: 258



Date: 16-09-2003


Day number: 259



Date: 17-09-2003


# Lerwick

Day number: 260



Date: 18-09-2003


Day number: 261



Date: 19-09-2003


# Lerwick

Day number: 262



Date: 20-09-2003


Day number: 263



Date: 21-09-2003


# Lerwick

Day number: 264



Date: 22-09-2003

Day number: 265






Date: 23-09-2003


# Lerwick

Day number: 266



Date: 24-09-2003


Day number: 267



Date: 25-09-2003


# Lerwick

Day number: 268



Date: 26-09-2003


Day number: 269



Date: 27-09-2003


# Lerwick

Day number: 270



Date: 28-09-2003


Day number: 271



Date: 29-09-2003


# Lerwick

Day number: 272




Date: 30-09-2003

Day number: 273




# Lerwick Observatory: Declination (degrees)




Hourly Mean Values Plotted By Bartels Solar Rotation Number

# Lerwick Observatory: Horizontal Intensity (nT)




Hourly Mean Values Plotted By Bartels Solar Rotation Number

# Lerwick Observatory: Vertical Intensity (nT)



Hourly Mean Values Plotted By Bartels Solar Rotation Number

# Lerwick 2003





### Monthly Mean Values for Lerwick Observatory 2003

Month	<i>D</i>	<i>H</i>	<i>I</i>	<i>X</i>	<i>Y</i>	<i>Z</i>	<i>F</i>	Data
January	-4° 20.3'	14936 nT	72° 49.3'	14893 nT	-1130 nT	48313 nT	50569 nT	<b>P</b>
February	-4° 18.9'	14935 nT	72° 49.4'	14892 nT	-1124 nT	48317 nT	50572 nT	<b>P</b>
March	-4° 17.3'	14937 nT	72° 49.2'	14895 nT	-1117 nT	48311 nT	50568 nT	<b>P</b>
April	-4° 16.9'	14939 nT	72° 49.0'	14897 nT	-1115 nT	48312 nT	50569 nT	<b>P</b>
May	-4° 15.5'	14940 nT	72° 49.0'	14899 nT	-1109 nT	48314 nT	50571 nT	<b>P</b>
June	-4° 15.2'	14947 nT	72° 48.7'	14906 nT	-1109 nT	48321 nT	50580 nT	<b>P</b>
July	-4° 13.9'	14948 nT	72° 48.7'	14907 nT	-1103 nT	48324 nT	50583 nT	<b>P</b>
August	-4° 12.8'	14943 nT	72° 49.1'	14903 nT	-1098 nT	48330 nT	50587 nT	<b>P</b>
September	-4° 12.2'	14942 nT	72° 49.2'	14902 nT	-1095 nT	48331 nT	50588 nT	<b>P</b>

## INDICES OF GEOMAGNETIC ACTIVITY

Lerwick Observatory

September 2003

Day	K - INDICES FOR THREE-HOUR INTERVAL								SUM
	00-03	03-06	06-09	09-12	12-15	15-18	18-21	21-24	
1	2	2	2	1	1	3	4	1	16
2	3	1	1	2	3	2	1	3	16
3	2	2	2	2	2	3	2	3	18
4	3	3	3	2	3	3	3	4	24
5	4	2	3	2	3	3	1	1	19
6	2	1	1	1	2	2	2	2	13
7	2	1	0	0	1	2	1	0	7
8	1	0	0	0	2	1	1	3	8
9	2	1	1	3	3	3	3	4	20
10	5	1	2	3	3	3	3	3	23
11	4	3	1	2	2	3	3	3	21
12	2	2	2	1	2	2	2	0	13
13	3	2	1	1	0	1	1	2	11
14	1	0	0	0	1	1	1	2	6
15	1	0	0	0	0	1	2	3	7
16	3	4	3	2	5	3	5	3	28
17	5	4	3	3	6	5	4	5	35
18	5	4	3	4	4	5	5	5	35
19	4	3	4	4	4	4	4	3	30
20	3	4	3	2	3	4	2	3	24
21	3	2	2	2	3	3	3	2	20
22	3	2	2	2	3	3	4	0	19
23	2	3	1	1	3	3	3	2	18
24	6	4	4	2	4	3	4	4	31
25	5	3	3	3	3	3	4	3	27
26	2	2	2	2	2	3	3	2	18
27	3	1	1	0	0	0	0	3	8
28	1	0	1	1	1	2	2	0	8
29	1	1	0	1	1	1	1	2	8
30	0	0	1	0	1	2	1	2	7

Lower bound (nT) for the range for each index value at Lerwick Observatory									
K-Index									
0	1	2	3	4	5	6	7	8	9
0	10	20	40	80	140	240	400	660	1000

### The aa Index

Date	Day	K-North	K-South	(a)	(b)	(c)	(d)	(e)
01-09-03	244	2 3 3 1 1 3 4 2	1 3 3 2 1 2 3 2	25	20	22	23	23
02-09-03	245	3 2 2 2 3 2 2 3	2 1 1 2 4 2 1 2	22	19	16	24	20
03-09-03	246	3 3 2 3 3 3 3 3	2 3 3 4 2 2 3 2	30	28	32	26	29
04-09-03	247	3 3 3 2 4 3 4 4	2 3 4 4 4 3 4 4	40	47	35	52	44
05-09-03	248	4 3 3 3 3 3 2 1	3 4 3 3 3 3 2 2	30	32	39	23	31
06-09-03	249	3 2 2 2 2 2 2 3	1 2 1 3 2 2 1 1	20	14	18	16	17
07-09-03	250	2 1 1 0 1 1 1 1	1 1 0 0 0 2 1 0	8	6	7	8	7
08-09-03	251	1 0 0 1 2 2 2 3	0 0 0 1 1 1 1 3	12	9	4	17	11
09-09-03	252	2 1 2 4 4 3 4 4	1 2 3 4 4 3 4 4	38	41	27	52	39
10-09-03	253	5 2 2 4 3 3 3 3	4 2 2 4 3 3 3 2	40	33	43	30	36
11-09-03	254	4 3 1 2 2 3 4 3	3 3 2 3 2 2 2 2	31	22	28	25	27
12-09-03	255	2 3 3 2 2 2 2 1	2 3 2 2 3 1 1 1	19	17	22	14	18
13-09-03	256	3 3 2 1 1 0 1 3	2 3 3 2 0 1 2 2	17	18	23	12	17
14-09-03	257	1 1 1 0 1 1 1 3	1 1 2 0 1 1 0 1	10	8	8	10	9
15-09-03	258	2 0 0 0 0 1 2 4	0 0 0 0 0 0 3 3	13	10	4	19	12
16-09-03	259	4 3 4 4 4 3 4 3	3 3 4 4 6 5 4 2	48	67	49	67	58
17-09-03	260	5 5 4 5 5 6 5 5	3 4 4 5 6 4 4 4	104	76	77	103	90
18-09-03	261	5 4 4 5 4 5 4 5	4 4 4 5 4 4 4 4	79	65	75	70	72
19-09-03	262	4 3 4 4 4 4 4 3	3 3 5 5 5 3 3 2	52	57	60	49	54
20-09-03	263	4 3 4 3 3 4 2 3	2 3 4 5 4 4 2 1	40	44	49	35	42
21-09-03	264	4 2 2 3 3 4 3 3	4 2 3 3 4 3 2 2	34	33	33	35	34
22-09-03	265	3 3 3 4 3 3 4 1	2 3 3 4 3 3 3 1	35	31	37	29	33
23-09-03	266	2 4 2 2 3 4 3 3	2 3 2 3 3 3 2 3	32	26	25	33	29
24-09-03	267	5 4 4 3 3 3 5 4	3 4 5 4 4 3 3 3	59	51	63	47	55
25-09-03	268	4 3 3 4 3 4 5 4	3 3 4 5 4 3 3 2	54	46	51	49	50
26-09-03	269	3 2 3 2 2 3 4 3	2 2 4 4 3 2 2 2	29	29	31	27	29
27-09-03	270	3 1 1 1 0 0 1 3	2 1 0 0 0 0 1 1	12	6	11	8	9
28-09-03	271	1 0 1 1 1 2 2 1	0 0 0 2 1 2 0 0	9	7	6	10	8
29-09-03	272	1 2 0 1 1 1 1 2	0 0 0 2 0 0 0 1 Est	9	5 Est	7 Est	7 Est	7 Est
30-09-03	273	0 0 1 1 1 2 1 2	2 2 2 0 2 2 0 2 Est	8	13 Est	9 Est	12 Est	11 Est
Monthly mean value =				30.7 Est				

- (a) The northern daily mean value,  $Aa_n$
- (b) The southern daily mean value,  $Aa_s$
- (c) The mean value of aa for the interval 00-12 UT
- (d) The mean value of aa for the interval 12-24 UT
- (e) The daily mean value of aa ( $Aa$ )

#### Notes

- i. The values are rounded to the nearest integer.
- iii. The units of the aa index are nT.
- iv. The values shown here are provisional. The definitive values are computed and published by the International Service for Geomagnetic Indices, Paris
- iv. Est indicates values that are not definitive.

## LERWICK RAPID VARIATIONS

### SI and SSCs

Date	Time (UT)	Type	Quality	H (nT)	D (min)	Z (nT)
NONE						

### SFEs

Date	Start (UT)	Maximum (UT)	End (UT)	H (nT)	D (min)	Z (nT)
NONE						

### Notes

- i. For SIs and SSCs a \* indicates that the principal impulse was preceded by a smaller reversed impulse.
- ii. The quality of SIs and SSCs are classified as follows:  
A = very distinct  
B = fair, ordinary, but unmistakable  
C = doubtful
- iii. The amplitudes given are for the first chief movement of the event.